



Opportunities and limits of the one gene approach: the ability of *Atoh1* to differentiate and maintain hair cells depends on the molecular context

Israt Jahan, Ning Pan and Bernd Fritzscht*

Department of Biology, University of Iowa, Iowa City, IA, USA

Edited by:

Andy Groves, Baylor College of Medicine, USA

Reviewed by:

Thomas Schimmang, Consejo Superior de Investigaciones Científicas, Spain
Albert Edge, Harvard Medical School, USA

*Correspondence:

Bernd Fritzscht, Department of Biology, College of Liberal Arts and Sciences, University of Iowa, 143 Biology Building, 129 E Jefferson Street, Iowa City, IA 52242-1324, USA
e-mail: bernd-fritzscht@uiowa.edu

Atoh1 (*Math1*) was the first gene discovered in ear development that showed no hair cell (HC) differentiation when absent and could induce HC differentiation when misexpressed. These data implied that *Atoh1* was both necessary and sufficient for hair cell development. However, other gene mutations also result in loss of initially forming HCs, notably null mutants for *Pou4f3*, *Barhl1*, and *Gfi1*. HC development and maintenance also depend on the expression of other genes (*Sox2*, *Eya1*, *Gata3*, *Pax2*) and several genes have been identified that can induce HCs when misexpressed (*Jag1*) or knocked out (*Lmo4*). In the ear *Atoh1* is not only expressed in HCs but also in some supporting cells and neurons that do not differentiate into HCs. Simple removal of one gene, *Neurod1*, can de-repress *Atoh1* and turns those neurons into HCs suggesting that *Neurod1* blocks *Atoh1* function in neurons. *Atoh1* expression in inner pillar cells may also be blocked by too many *Hes/Hey* factors but conversion into HCs has only partially been achieved through *Hes/Hey* removal. Detailed analysis of cell cycle exit confirmed an apex to base cell cycle exit progression of HCs of the organ of Corti. In contrast, *Atoh1* expression progresses from the base toward the apex with a variable delay relative to the cell cycle exit. Most HCs exit the cell cycle and are thus defined as precursors before *Atoh1* is expressed. *Atoh1* is a potent differentiation factor but can differentiate and maintain HCs only in the ear and when other factors are co-expressed. Upstream factors are essential to regulate *Atoh1* level of expression duration while downstream, co-activated by other factors, will define the context of *Atoh1* action. We suggest that these insights need to be taken into consideration and approaches beyond the simple *Atoh1* expression need to be designed able to generate the radial and longitudinal variations in hair cell types for normal function of the organ of Corti.

Keywords: *Atoh1*, hair cells, development, regeneration

INTRODUCTION

The idea that single genes might be responsible for hair cell (HC) development and thus could be used to regenerate HCs and restore hearing was born in the late 1990s: Mice with a deletion of the Pou domain gene *Pou4f3* (aka *Brn3c*, *Brn3.1*) were completely deaf, “owing to a failure of HCs to appear in the inner ear, with subsequent loss of cochlear and vestibular ganglia” (Erkman et al., 1996). This mouse mutant derived conclusion was soon followed by data on human mutations showing that a truncating mutation of the human *POU4f3* gene is the basis of DFNA15, resulting in progressive hearing loss (Vahava et al., 1998). Subsequent work showed that HCs initially form and develop normal in *Pou4f3* mutants, but eventually die in a base to apex progression (Xiang et al., 2003; Hertzano et al., 2004). While the initial work claimed loss of all sensory neurons, later work showed that some neurons remain for 6 months in a dedifferentiated organ of Corti (OC) that shows *Atoh1-lacZ* and *Myo7a* positive cells (Pauley et al., 2008). The original claim of “failure of HCs to appear” was thus transformed into a rather normal initial development followed by HC death. *Pou4f3* is now recognized as a maintenance factor

for HCs, like *Gfi1* and *Barhl1* (Li et al., 2002; Hertzano et al., 2004) that is expressed in adult HCs through complex regulation, including possibly the bHLH gene *Atoh1* (Ahmed et al., 2012; Masuda et al., 2012).

Why is this background information on *Pou4f3* relevant for the discussion of the role of *Atoh1* (aka *Math1*) for HC differentiation and maintenance? In the following we will explore that *Atoh1* has much in common with *Pou4f3* in terms of claims raised as a gene that is “necessary and sufficient” for HC differentiation (Chen et al., 2002; Giraldez and Fritzscht, 2007; Groves et al., 2013). In contrast to this claim, the millions of neurons outside the ear expressing *Atoh1* (Mulvaney and Dabdoub, 2012) never turn into HCs, suggesting that *Atoh1* is not sufficient to induce HCs everywhere where *Atoh1* is expressed. Only the molecularly unclear context of certain cells of the ear allows *Atoh1* to drive HC differentiation and maintenance. Even in the ear, *Atoh1* is expressed in many cells (Matei et al., 2005) that require additional manipulations to turn into HCs (Jahan et al., 2010), indicating that expression of *Atoh1* in the ear does not guarantee differentiation of all cells into HCs. As with *Pou4f3*,

it appears that *Atoh1* absence is compatible with some cellular differentiation, indicating that *Atoh1* is not defining HCs but is differentiating them (Jahan et al., 2012). The delayed and profound loss of HCs in a “self-terminating” *Atoh1* system (Pan et al., 2012) and hypomorphic *Atoh1* mutant (Sheykhosslami et al., 2013) suggests an essential role in maintenance, possibly including adult expression of *Pou4f3* (Masuda et al., 2012). Consistent with *Atoh1* being an essential differentiation and maintenance factor for HC is the fact that overexpression can rescue HCs (Yang et al., 2012). Like *Pou4f3*, *Atoh1* is necessary to differentiate and maintain HCs. It remains to be shown whether forced expression of *Atoh1* (Kelly et al., 2012) can differentiate HCs when certain factors are absent (Zou et al., 2004; Kiernan et al., 2005; Bouchard et al., 2010; Ahmed et al., 2012; Duncan and Fritzsche, 2013; Schimmang, 2013) that define the context for *Atoh1* action in the ear thus providing the competency to respond to *Atoh1* protein. Below we explore some issues related to *Atoh1* function that remain underexplored in many contemporary reviews and propose novel strategies to maintain HCs.

EXPRESSION OF *Atoh1* OUTSIDE THE EAR DOES NOT LEAD TO HC DIFFERENTIATION


Atoh1 was isolated from cerebellar granule cells, the largest population of neurons in the human brain, amounting to over 60 billion neurons (Ben-Arie et al., 1997; Herculano-Houzel, 2009). *Atoh1* is expressed in the proliferative precursor population of the external granule cell layer where it is needed to generate the billions of granule cells (Pan et al., 2009). *Atoh1* is also essential for medulloblastoma progression and *Atoh1* removal reduces the progression of this childhood tumor (Flora et al., 2009). In contrast to this expression of *Atoh1* in proliferating precursors in the CNS, the expression of *Atoh1* in the mouse cochlea is predominantly in post-mitotic HCs, with a possible overlap of *Atoh1* expression and cell cycle exit in the basal turn HCs (Ruben, 1967; Matei et al., 2005; Lee et al., 2006). A pulse-chase experiment using BrdU or EdU labeling followed by *in situ* hybridization for *Atoh1* around E14 is needed to verify this suggestion of possible *Atoh1* expression in proliferating HC precursors. In the apex there is no expression of *Atoh1* prior to cell cycle exit, indicating that HC precursor specification and cell cycle exit is independent of *Atoh1* (Jahan et al., 2013; Kopecky et al., 2013). Both premature expression of *Atoh1* in *Neurod1* null mutants (Jahan et al., 2010) or delayed expression of *Atoh1* in *Lmx1a* null mice (Nichols et al., 2008) results in aberrant development of HCs, implying that onset and level of expression of *Atoh1* is tightly regulated to ensure normal differentiation of the right HC type at the right place (Jahan et al., 2013). Importantly, forced expression of *Atoh1* can in postnatal mice induce supporting cell conversion (Liu et al., 2014a) and induces proliferation (Kelly et al., 2012), showing that under these forced conditions *Atoh1* exerts functions beyond its tightly regulated function in the embryonic ear. In summary, one of the conditions in which *Atoh1* expression in the ear differs from other systems is its expression presumably exclusively in post-mitotic undifferentiated HC precursors whereas in other developing systems *Atoh1* is primarily expressed in proliferating precursors.

UPSTREAM AND DOWNSTREAM INTERACTIONS OF *Atoh1*

Before *Atoh1* can differentiate post-mitotic HC precursors into HCs, the HC precursors have to be specified in the right place and have to receive a signal to exit the cell cycle. Numerous TFs have been identified that are expressed prior to *Atoh1* and affect HC differentiation. For example, *Sox2* hypomorphic mice (Kiernan et al., 2005), *Pax2* null mice (Bouchard et al., 2010), *Eya1* null mice (Zou et al., 2004), and *Gata3* conditional null mice (Duncan and Fritzsche, 2013) all show no differentiation of HCs in the cochlea duct but may show variable development of some vestibular HCs, suggesting a unique combinatorial requirement of these genes for cochlear HC development. Misexpression of *Jag1* (Pan et al., 2010) or *Sox2* (Pan et al., 2013) as well as loss of *Lmo4* (Deng et al., 2014) can induce ectopic formation of HCs. In particular work on *Eya1/Six1* showed that *Atoh1* is but an essential link in a succession of decision making steps (Ahmed et al., 2012) toward HC differentiation (Figure 1) with unknown regulatory complexity.

Atoh1 regulates the expression of hundreds of downstream genes (Klisch et al., 2011). Some of these genes are TFs that in turn regulate expression of several hundred downstream genes. One of the TFs that are regulated by *Atoh1*, is *Neurod1*. *Atoh1* is in a positive autoregulatory loop whereby *Atoh1* stimulates its own expression through an enhancer sequence (Figure 1). Such loops are typically counterbalanced by negative feedback to ensure upper limits of expression. *Neurod1* is part of this negative feedback loop and controls the level of *Atoh1* expression in developing systems such as the cerebellum (Pan et al., 2009), the intestine (Itkin-Ansari et al., 2005), and the ear (Jahan et al., 2010, 2013). Absence of *Neurod1* causes prolonged expression of *Atoh1* in precursor cells (external granule cell layer) of the cerebellum that are unable to migrate and differentiate and eventually die (Pan et al., 2009). In the ear, absence of *Neurod1* causes transformation of sensory neurons into HCs through disinhibition of a transient *Atoh1* in neurons (Jahan et al., 2010) and disruption of the patterning of the OC by altering the HC and supporting cell types (Jahan et al., 2013). Some regulation of *Atoh1* is also reported in mutants of *Hes1/5* (Zine et al., 2001, 2014) and *Hey1/2* (Benito-Gonzalez and Doetzlhofer, 2014) but results only in additional formation of HCs outside the OC with limited effects on the patterning of HCs and supporting cells in the OC. *Atoh1* is not only regulating the expression of downstream genes but also suppresses upstream genes such as *Sox2* (Figure 1). In fact, downregulation of *Sox2* appears to be a crucial step for the transition from HC precursors to differentiated HCs (Dabdoub et al., 2008) in agreement with many other differentiating neurosensory system (Reiprich and Wegner, 2014).

Combined, these data show that the early implications of *Atoh1* as the “sole” factor necessary and sufficient to make HCs have to be adjusted to accommodate the emerging concept of *Atoh1* integration into a gene network that allows a coordinated transition from the placodal stage to the fully differentiated HC (Ahmed et al., 2012). Arguably, *Atoh1* is enabling a very essential step in this progression toward a HC, but is apparently not needed for precursors to exit the cell cycle and to initiate HC differentiation (Jahan et al., 2012). However, *Atoh1* is a key to HC differentiation (Kelly et al., 2012) and its continued expression



Using the same LacZ knockin model as previous papers (Bermingham et al., 1999; Woods et al., 2004), a follow up paper on homozygotic Atoh1-LacZ mice showed continued presence of a single row of undifferentiated LacZ positive cells (Fritzsch et al., 2005) which were spared by the otherwise prevalent apoptosis of most HC precursors (Chen et al., 2002). Subsequent work demonstrated that fluorescent GFP marker (Chen et al., 2002) appeared in nearly every inner pillar cell (Matei et al., 2005;

Fritzsche et al., 2015b). Furthermore, a novel mouse line using the same enhancer element to drive Cre showed similar expression of *Atoh1* in many inner pillar cells (Matei et al., 2005). These data implied, but did not prove beyond doubt that *Atoh1* was expressed in inner pillar cells and inferred that the remaining *Atoh1*-LacZ positive cells in mutants were indeed supporting cells (possibly inner pillar cells) as originally claimed (Bermingham et al., 1999). Further work using a conditional approach to eliminate *Atoh1* resulted in nearly identical data, implying that the surviving cells in the absence of *Atoh1* might indeed be inner pillar cells in the OC (Pan et al., 2011). Additional work has meanwhile confirmed with different techniques that *Atoh1* is indeed prominently expressed in inner pillar cells (Driver et al., 2013). *Atoh1* expression in inner pillar cells may be counterbalanced by Hes and Hey factors (Doetzlhofer et al., 2009) and a subsequent paper showed occasional conversion of inner pillar cells to HCs (Benito-Gonzalez and Doetzlhofer, 2014). *Atoh1* expression has also been reported in delaminating sensory neurons (Matei et al., 2005) and elimination of *Neurod1* suffices to turn some neurons into HCs expressing *Atoh1* and *Myo7a* (Jahan et al., 2010). Combined, these data suggest that *Atoh1* expression alone does not suffice to turn just any cell in the ear into a HC as co-expressed factors may inhibit this. At least inner pillar cells may be able to survive without *Atoh1* protein while maintaining LacZ expression of the *Atoh1* locus (Matei et al., 2005; Driver et al., 2013) and are not transformed to HCs even under forced ubiquitous expression of *Atoh1* (Kelly et al., 2012).

More recent data provide yet a more complicated picture of lack of *Atoh1* expression on HC and OC differentiation. Using an *Atoh1* enhancer to drive Cre that activates the Cre only upon presence of *Atoh1* protein combined with floxed *Atoh1* generates a “self-terminating” system that results in loss of *Atoh1* after a transient presence of *Atoh1* protein (Pan et al., 2012). The level of *Atoh1* protein depends on the speed with which the Cre can excise the floxed *Atoh1* and how long residual *Atoh1* protein remains in the cell. Thus, while all cells will see recombination of the LoxP flanked *Atoh1*, this varies between HCs and thus results in different delay lines of HC precursor apoptosis (Pan et al., 2012). While many HC precursors die rapidly, others survive for several days. Moreover, stretches of the first row of outer HCs survive adjacent to well differentiated inner pillar cells indicating an unusual difference in susceptibility between inner and outer HCs as well as within HC rows in a base to apical gradient. This conclusion is also supported by transgenic knockin mouse where *Atoh1* is replaced by *Neurog1* (Jahan et al., 2012) which shows that some HC precursors can survive without ever expressing *Atoh1*. A recently available hypomorph mutant of *Atoh1* shows a somewhat similar picture of longitudinal and a less clear radial HC loss (Sheykholslami et al., 2013) indicating that *Atoh1* needs to be present at a critical level to assure long term HC viability.

Data using inducible Cre expression have complicated this picture even further by showing a rapid and complete loss of all HCs when Cre is induced at different stages of late development (Cai et al., 2013; Chonko et al., 2013). Some claims about abortive transdifferentiation of supporting cells into HCs (Cai et al., 2013) need to be considered in the context of *Atoh1* expression in one specific type of supporting cell, the inner pillar cell (Matei et al.,

2005; Driver et al., 2013; Fritzsche et al., 2015b). Despite these minor discrepancies, all papers confirm earlier work and demonstrate that *Atoh1* expression is needed to mature and maintain HCs.

In summary, *Atoh1* is, much like *Pou4f3*, a critical factor for HC differentiation and long term maintenance. *Atoh1* is involved in regulating *Pou4f3* whereas its long term expression may be dependent on *Atoh1* expression. Further work combining the recently reported hypomorphic allele (Sheykholslami et al., 2013) with conditional deletion of a floxed *Atoh1* allele (Pan et al., 2012) could detail how level of *Atoh1* expression and duration combine for normal HC maturation and maintenance.

SUMMARY AND OUTLOOK

Why is it important to go beyond the idea of “necessary and sufficient” for *Atoh1* function in the ear? First, while unregulated expression of *Atoh1* can convert most ear cells into hair cells (Kelly et al., 2012), nobody has been able to regenerate the two types of HCs that are essential for normal OC function in the right proportion and the right distribution to ensure function (Beurg et al., 2014). In fact, our limited insights into the molecular basis of this crucial aspect of HC differentiation (Jahan et al., 2013) are not profound enough to regenerate the right type of HC (Liu et al., 2014b) to ensure normal function. Defining the molecular context needed for HC type specific differentiation in conjunction with defined levels of *Atoh1* expression (Jahan et al., 2013) and controlled changes of *Atoh1* expression over time (Ahmed et al., 2012) will be needed to move forward.

Second, most HCs generated with *Atoh1* treatment alone have limited long term viability. In part this may relate to the progressive loss of *Atoh1* in these experiments that may be needed to maintain long term *Pou4f3* expression (Masuda et al., 2012), but in part it may also relate to an unstable transformation into HC that requires recapitulating the specification sequence of HCs precursors and their differentiation. Such critical steps might include expression of additional factors prior to and in addition to *Atoh1* or the prolonged expression of critical levels of *Atoh1*. Human hearing loss may show partial dedifferentiation of the OC with profound local differences comparable to experimental animals (Taylor et al., 2012). A “one size fits all” approach to such heterogeneity may result in incomplete restoration.

Finally, while the single gene approach to HC regeneration has been extremely influential to catapult much research forward, it is now time to reflect why this approach has not lived up to its promise. We therefore suggest more complex procedures that recapitulate steps in development of the OC in addition to *Atoh1*. For example, expressing *Eya1*, *Pax2*, *Sox2*, *Jag1*, *Foxg1*, *Neurod1*, *Neurog1*, and *Gata3* prior to *Atoh1* expression may “prime” remaining cells of the OC to respond to *Atoh1*. Alternatively, combining *Atoh1* with downstream essential genes for HC maintenance that are only partially regulated by *Atoh1* (Ahmed et al., 2012), such as *Pou4f3*, could define the context for HC differentiation. Moreover, using transient expression of *Atoh1* in already differentiated HCs might prolong their viability (Yang et al., 2012), possibly long enough to sidestep the need for OC regeneration in elderly people suffering from early stages of neurosensory hearing loss. Given the projected massive occurrence of hearing

loss in the next 25 years, ideas revolving around maintenance of HCs using Atoh1 alone might provide more-short term benefit compared to currently impossible reconstitution of the OC after long term HC loss. Given the ability of Atoh1 to transdifferentiate supporting cells in certain conditions (Liu et al., 2014a), it might be necessary to replace Atoh1 by other bHLH genes that can accomplish long term maintenance of HCs without risk of transforming supporting cells into HCs. We are currently working on such approaches using novel mouse models to differentiate HCs in the absence or at most transient presence of Atoh1.

ACKNOWLEDGMENTS

This work was supported by NIH (P30 DC 010362, R03 DC013655), NASA Base Program and the OVPR, University of Iowa.

REFERENCES

- Ahmed, M., Wong, E. Y., Sun, J., Xu, J., Wang, F., and Xu, P.-X. (2012). Eya1-Six1 interaction is sufficient to induce hair cell fate in the cochlea by activating Atoh1 expression in cooperation with Sox2. *Dev. Cell* 22, 377–390. doi: 10.1016/j.devcel.2011.12.006
- Ben-Arie, N., Bellen, H. J., Armstrong, D. L., McCall, A. E., Gordadze, P. R., Guo, Q., et al. (1997). Math1 is essential for genesis of cerebellar granule neurons. *Nature* 390, 169–172. doi: 10.1038/36579
- Benito-Gonzalez, A., and Doetzlhofer, A. (2014). Hey1 and Hey2 Control the Spatial and Temporal pattern of mammalian auditory hair cell differentiation downstream of hedgehog signaling. *J. Neurosci.* 34, 12865–12876. doi: 10.1523/JNEUROSCI.1494-14.2014
- Bermingham, N. A., Hassan, B. A., Price, S. D., Vollrath, M. A., Ben-Arie, N., Eatock, R. A., et al. (1999). Math1: an essential gene for the generation of inner ear hair cells. *Science* 284, 1837–1841. doi: 10.1126/science.284.5421.1837
- Beurg, M., Xiong, W., Zhao, B., Müller, U., and Fettiplace, R. (2014). Subunit determination of the conductance of hair-cell mechanotransducer channels. *Proc. Natl. Acad. Sci. U.S.A.* doi: 10.1073/pnas.1420906112. [Epub ahead of print].
- Bouchard, M., de Caprona, D., Busslinger, M., Xu, P., and Fritzsche, B. (2010). Pax2 and Pax8 cooperate in mouse inner ear morphogenesis and innervation. *BMC Dev. Biol.* 10:89. doi: 10.1186/1471-213X-10-89
- Cai, T., Seymour, M. L., Zhang, H., Pereira, F. A., and Groves, A. K. (2013). Conditional deletion of Atoh1 reveals distinct critical periods for survival and function of hair cells in the organ of Corti. *J. Neurosci.* 33, 10110–10122. doi: 10.1523/JNEUROSCI.5606-12.2013
- Chen, P., Johnson, J. E., Zoghbi, H. Y., and Segil, N. (2002). The role of Math1 in inner ear development: uncoupling the establishment of the sensory primordium from hair cell fate determination. *Development* 129, 2495–2505.
- Chonko, K. T., Jahan, I., Stone, J., Wright, M. C., Fujiyama, T., Hoshino, M., et al. (2013). Atoh1 directs hair cell differentiation and survival in the late embryonic mouse inner ear. *Dev. Biol.* 381, 401–410. doi: 10.1016/j.ydbio.2013.06.022
- Dabdoub, A., Puligilla, C., Jones, J. M., Fritzsche, B., Cheah, K. S., Pevny, L. H., et al. (2008). Sox2 signaling in prosensory domain specification and subsequent hair cell differentiation in the developing cochlea. *Proc. Natl. Acad. Sci. U.S.A.* 105, 18396–18401. doi: 10.1073/pnas.0808175105
- Deng, M., Luo, X. J., Pan, L., Yang, H., Xie, X., Liang, G., et al. (2014). LMO4 functions as a negative regulator of sensory organ formation in the mammalian cochlea. *J. Neurosci.* 34, 10072–10077. doi: 10.1523/JNEUROSCI.0352-14.2014
- Doetzlhofer, A., Basch, M. L., Ohshima, T., Gessler, M., Groves, A. K., and Segil, N. (2009). Hey2 regulation by FGF provides a Notch-independent mechanism for maintaining pillar cell fate in the organ of Corti. *Dev. Cell* 16, 58–69. doi: 10.1016/j.devcel.2008.11.008
- Domínguez-Frutos, E., López-Hernández, I., Vendrell, V., Neves, J., Gallozzi, M., Gutsche, K., et al. (2011). N-myc controls proliferation, morphogenesis, and patterning of the inner ear. *J. Neurosci.* 31, 7178–7189. doi: 10.1523/JNEUROSCI.0785-11.2011
- Driver, E. C., Sillers, L., Coate, T. M., Rose, M. F., and Kelley, M. W. (2013). The Atoh1-lineage gives rise to hair cells and supporting cells within the mammalian cochlea. *Dev. Biol.* 376, 86–98. doi: 10.1016/j.ydbio.2013.01.005
- Duncan, J. S., and Fritzsche, B. (2013). Continued expression of GATA3 is necessary for cochlear neurosensory development. *PLoS ONE* 8:e62046. doi: 10.1371/journal.pone.0062046
- Erkman, L., McEvilly, R. J., Luo, L., Ryan, A. K., Hooshmand, F., O'Connell, S. M., et al. (1996). Role of transcription factors Brn-3.1 and Brn-3.2 in auditory and visual system development. *Nature* 381, 603–606. doi: 10.1038/381603a0
- Flora, A., Klisch, T. J., Schuster, G., and Zoghbi, H. Y. (2009). Deletion of Atoh1 disrupts Sonic Hedgehog signaling in the developing cerebellum and prevents medulloblastoma. *Science* 326, 1424–1427. doi: 10.1126/science.1181453
- Fritzsche, B., Beisel, K. W., and Hansen, L. A. (2006). The molecular basis of neurosensory cell formation in ear development: a blueprint for hair cell and sensory neuron regeneration? *Bioessays* 28, 1181–1193. doi: 10.1002/bies.20502
- Fritzsche, B., Jahan, I., Pan, N., and Elliott, K. L. (2015a). Evolving gene regulatory networks into cellular networks guiding adaptive behavior: an outline how single cells could have evolved into a centralized neurosensory system. *Cell Tissue Res.* 359, 295–313. doi: 10.1007/s00441-014-2043-1
- Fritzsche, B., Matei, V. A., Nichols, D. H., Bermingham, N., Jones, K., Beisel, K. W., et al. (2005). Atoh1 null mice show directed afferent fiber growth to undifferentiated ear sensory epithelia followed by incomplete fiber retention. *Dev. Dyn.* 233, 570–583. doi: 10.1002/dvdy.20370
- Fritzsche, B., and Nichols, D. (1993). Dil reveals a prenatal arrival of efferents at the differentiating otocyst of mice. *Hear. Res.* 65, 51–60. doi: 10.1016/0378-5955(93)90200-K
- Fritzsche, B., Pan, N., Jahan, I., and Elliott, K. L. (2015b). Inner ear development: building a spiral ganglion and an organ of Corti out of unspecified ectoderm. *Cell Tissue Res.* doi: 10.1007/s00441-014-2031-5. [Epub ahead of print].
- Giraldez, F., and Fritzsche, B. (2007). The molecular biology of ear development—“Twenty years are nothing.” *Int. J. Dev. Biol.* 51, 429. doi: 10.1387/ijdb.072390fg
- Gokofski, K. K., Wu, H. H., Beites, C. L., Kim, J., Kim, E. J., Matzuk, M. M., et al. (2011). Activin and GDF11 collaborate in feedback control of neuroepithelial stem cell proliferation and fate. *Development* 138, 4131–4142. doi: 10.1242/dev.065870
- Groves, A. K., Zhang, K. D., and Fekete, D. M. (2013). The genetics of hair cell development and regeneration. *Ann. Rev. Neurosci.* 36, 361–381. doi: 10.1146/annurev-neuro-062012-170309
- Herculano-Houzel, S. (2009). The human brain in numbers: a linearly scaled-up primate brain. *Front. Hum. Neurosci.* 3:31. doi: 10.3389/neuro.09.031
- Hertzano, R., Montcouquiol, M., Rashi-Elkeles, S., Elkou, R., Yucel, R., Frankel, W. N., et al. (2004). Transcription profiling of inner ears from Pou4f3 (ddl/ddl) identifies Gfi1 as a target of the Pou4f3 deafness gene. *Hum. Mol. Genet.* 13, 2143–2153. doi: 10.1093/hmg/ddh218
- Imayoshi, I., and Kageyama, R. (2014). bHLH factors in self-renewal, multipotency, and fate choice of neural progenitor cells. *Neuron* 82, 9–23. doi: 10.1016/j.neuron.2014.03.018
- Itkin-Ansari, P., Marcora, E., Geron, I., Tyrberg, B., Demeterco, C., Hao, E., et al. (2005). NeuroD1 in the endocrine pancreas: localization and dual function as an activator and repressor. *Dev. Dyn.* 233, 946–953. doi: 10.1002/dvdy.20443
- Jahan, I., Pan, N., Kersigo, J., Calisto, L. E., Morris, K. A., Kopecky, B., et al. (2012). Expression of Neurog1 instead of Atoh1 can partially rescue organ of Corti cell survival. *PLoS ONE* 7:e30853. doi: 10.1371/journal.pone.0030853
- Jahan, I., Pan, N., Kersigo, J., and Fritzsche, B. (2010). Neurod1 suppresses hair cell differentiation in ear ganglia and regulates hair cell subtype development in the cochlea. *PLoS ONE* 5:e11661. doi: 10.1371/journal.pone.0011661
- Jahan, I., Pan, N., Kersigo, J., and Fritzsche, B. (2013). Beyond generalized hair cells: molecular cues for hair cell types. *Hear. Res.* 297, 30–41. doi: 10.1016/j.heares.2012.11.008
- Kelly, M. C., Chang, Q., Pan, A., Lin, X., and Chen, P. (2012). Atoh1 directs the formation of sensory mosaics and induces cell proliferation in the postnatal mammalian cochlea in vivo. *J. Neurosci.* 32, 6699–6710. doi: 10.1523/JNEUROSCI.5420-11.2012
- Kiernan, A. E., Pelling, A. L., Leung, K. K., Tang, A. S., Bell, D. M., Tease, C., et al. (2005). Sox2 is required for sensory organ development in the mammalian inner ear. *Nature* 434, 1031–1035. doi: 10.1038/nature03487
- Klisch, T. J., Xi, Y., Flora, A., Wang, L., Li, W., and Zoghbi, H. Y. (2011). In vivo Atoh1 targetome reveals how a proneural transcription factor regulates cerebellar development. *Proc. Natl. Acad. Sci. U.S.A.* 108, 3288–3293. doi: 10.1073/pnas.1100230108

- Kopecky, B., DeCook, R., and Fritzsche, B. (2012). Mutational ataxia resulting from abnormal vestibular acquisition and processing is partially compensated for. *Behav. Neurosci.* 126, 301. doi: 10.1037/a0026896
- Kopecky, B. J., Jahan, I., and Fritzsche, B. (2013). Correct timing of proliferation and differentiation is necessary for normal inner ear development and auditory hair cell viability. *Dev. Dyn.* 242, 132–147. doi: 10.1002/dvdy.23910
- Kopecky, B., Santi, P., Johnson, S., Schmitz, H., and Fritzsche, B. (2011). Conditional deletion of N-Myc disrupts neurosensory and non-sensory development of the ear. *Dev. Dyn.* 240, 1373–1390. doi: 10.1002/dvdy.22620
- Lee, Y.-S., Liu, F., and Segil, N. (2006). A morphogenetic wave of p27Kip1 transcription directs cell cycle exit during organ of Corti development. *Development* 133, 2817–2826. doi: 10.1242/dev.02453
- Li, S., Price, S. M., Cahill, H., Ryugo, D. K., Shen, M. M., and Xiang, M. (2002). Hearing loss caused by progressive degeneration of cochlear hair cells in mice deficient for the Barhl1 homeobox gene. *Development* 129, 3523–3532.
- Liu, H., Pecka, J. L., Zhang, Q., Soukup, G. A., Beisel, K. W., and He, D. Z. (2014b). Characterization of transcriptomes of cochlear inner and outer hair cells. *J. Neurosci.* 34, 11085–11095. doi: 10.1523/JNEUROSCI.1690-14.2014
- Liu, Z., Fang, J., Dearman, J., Zhang, L., and Zuo, J. (2014a). *In vivo* generation of immature inner hair cells in neonatal mouse cochlea by ectopic Atoh1 expression. *PLoS ONE* 9:e89377. doi: 10.1371/journal.pone.0089377
- Ma, Q., Chen, Z., del Barco Barrantes, I., de la Pompa, J. L., and Anderson, D. J. (1998). Neurogenin1 is essential for the determination of neuronal precursors for proximal cranial sensory ganglia. *Neuron* 20, 469–482. doi: 10.1016/S0896-6273(00)80988-5
- Ma, Q., Kintner, C., and Anderson, D. J. (1996). Identification of neurogenin, a vertebrate neuronal determination gene. *Cell* 87, 43–52. doi: 10.1016/S0092-8674(00)81321-5
- Mantela, J., Jiang, Z., Ylikoski, J., Fritzsche, B., Zacksenhaus, E., and Pirvola, U. (2005). The retinoblastoma gene pathway regulates the postmitotic state of hair cells of the mouse inner ear. *Development* 132, 2377–2388. doi: 10.1242/dev.01834
- Masuda, M., Pak, K., Chavez, E., and Ryan, A. F. (2012). TFE2 and GATA3 enhance induction of POU4F3 and myosin VIIa positive cells in nonsensory cochlear epithelium by ATOH1. *Dev. Biol.* 372, 68–80. doi: 10.1016/j.ydbio.2012.09.002
- Matei, V., Pauley, S., Kaing, S., Rowitch, D., Beisel, K. W., Morris, K., et al. (2005). Smaller inner ear sensory epithelia in Neurog 1 null mice are related to earlier hair cell cycle exit. *Dev. Dyn.* 234, 633–650. doi: 10.1002/dvdy.20551
- Mulvaney, J., and Dabdoub, A. (2012). Atoh1, an essential transcription factor in neurogenesis and intestinal and inner ear development: function, regulation, and context dependency. *J. Assoc. Res. Otolaryngol.* 13, 281–293. doi: 10.1007/s10162-012-0317-4
- Nichols, D. H., Pauley, S., Jahan, I., Beisel, K. W., Millen, K. J., and Fritzsche, B. (2008). Lmx1a is required for segregation of sensory epithelia and normal ear histogenesis and morphogenesis. *Cell Tissue Res.* 334, 339–358. doi: 10.1007/s00441-008-0709-2
- Ono, K., Kita, T., Sato, S., O'Neill, P., Mak, S.-S., Paschaki, M., et al. (2014). FGFR1-Frs2/3 signalling maintains sensory progenitors during inner ear hair cell formation. *PLoS Genet.* 10:e1004118. doi: 10.1371/journal.pgen.1004118
- Pan, N., Jahan, I., Kersigo, J., Duncan, J. S., Kopecky, B., and Fritzsche, B. (2012). A novel Atoh1 “self-terminating” mouse model reveals the necessity of proper Atoh1 level and duration for hair cell differentiation and viability. *PLoS ONE* 7:e30358. doi: 10.1371/journal.pone.0030358
- Pan, N., Jahan, I., Kersigo, J., Kopecky, B., Santi, P., Johnson, S., et al. (2011). Conditional deletion of Atoh1 using Pax2-Cre results in viable mice without differentiated cochlear hair cells that have lost most of the organ of Corti. *Hear. Res.* 275, 66–80. doi: 10.1016/j.heares.2010.12.002
- Pan, N., Jahan, I., Lee, J. E., and Fritzsche, B. (2009). Defects in the cerebella of conditional Neurod1 null mice correlate with effective Tg(Atoh1-cre) recombination and granule cell requirements for Neurod1 for differentiation. *Cell Tissue Res.* 337, 407–428. doi: 10.1007/s00441-009-0826-6
- Pan, W., Jin, Y., Chen, J., Rottier, R. J., Steel, K. P., and Kiernan, A. E. (2013). Ectopic expression of activated notch or SOX2 reveals similar and unique roles in the development of the sensory cell progenitors in the mammalian inner ear. *J. Neurosci.* 33, 16146–16157. doi: 10.1523/JNEUROSCI.3150-12.2013
- Pan, W., Jin, Y., Stanger, B., and Kiernan, A. E. (2010). Notch signaling is required for the generation of hair cells and supporting cells in the mammalian inner ear. *Proc. Natl. Acad. Sci. U.S.A.* 107, 15798–15803. doi: 10.1073/pnas.1003089107
- Pauley, S., Kopecky, B., Beisel, K., Soukup, G., and Fritzsche, B. (2008). Stem cells and molecular strategies to restore hearing. *Panminerva Med.* 50, 41–53.
- Reiprich, S., and Wegner, M. (2014). From CNS stem cells to neurons and glia: sox for everyone. *Cell Tissue Res.* 359, 111–124. doi: 10.1007/100441-014-1909-6
- Rocha-Sanchez, S. M., Scheetz, L. R., Contreras, M., Weston, M. D., Korte, M., McGee, J., et al. (2011). Mature mice lacking Rbl2/p130 gene have supernumerary inner ear hair cells and supporting cells. *J. Neurosci.* 31, 8883–8893. doi: 10.1523/JNEUROSCI.5821-10.2011
- Ruben, R. J. (1967). Development of the inner ear of the mouse: a radioautographic study of terminal mitoses. *Acta Otolaryngol. (Suppl. 220)*, 1–44.
- Schimmang, T. (2013). Transcription factors that control inner ear development and their potential for transdifferentiation and reprogramming. *Hear. Res.* 297, 84–90. doi: 10.1016/j.heares.2012.11.001
- Schimmang, T., and Pirvola, U. (2013). Coupling the cell cycle to development and regeneration of the inner ear. *Semin. Cell Dev. Biol.* 24, 507–513. doi: 10.1016/j.semcdb.2013.04.004
- Sheykholeslami, K., Thimmappa, V., Nava, C., Bai, X., Yu, H., Zheng, T., et al. (2013). A new mutation of the Atoh1 gene in mice with normal life span allows analysis of inner ear and cerebellar phenotype in aging. *PLoS ONE* 8:e79791. doi: 10.1371/journal.pone.0079791
- Taylor, R. R., Jagger, D. J., and Forge, A. (2012). Defining the cellular environment in the organ of Corti following extensive hair cell loss: a basis for future sensory cell replacement in the cochlea. *PLoS ONE* 7:e30577. doi: 10.1371/journal.pone.0030577
- Vahava, O., Morell, R., Lynch, E. D., Weiss, S., Kagan, M. E., Ahituv, N., et al. (1998). Mutation in transcription factor POU4F3 associated with inherited progressive hearing loss in humans. *Science* 279, 1950–1954. doi: 10.1126/science.279.5358.1950
- Woods, C., Montcouquiol, M., and Kelley, M. W. (2004). Math1 regulates development of the sensory epithelium in the mammalian cochlea. *Nat. Neurosci.* 7, 1310–1318. doi: 10.1038/nn1349
- Xiang, M., Maklad, A., Pirvola, U., and Fritzsche, B. (2003). Brn3c null mutant mice show long-term, incomplete retention of some afferent inner ear innervation. *BMC Neurosci.* 4:2. doi: 10.1186/1471-2202-4-2
- Yang, S. M., Chen, W., Guo, W. W., Jia, S., Sun, J. H., Liu, H. Z., et al. (2012). Regeneration of stereocilia of hair cells by forced Atoh1 expression in the adult mammalian cochlea. *PLoS ONE* 7:e46355. doi: 10.1371/journal.pone.0046355
- Zine, A., Aubert, A., Qiu, J., Therianos, S., Guillemot, F., Kageyama, R., et al. (2001). Hes1 and Hes5 activities are required for the normal development of the hair cells in the mammalian inner ear. *J. Neurosci.* 21, 4712–4720.
- Zine, A., Löwenheim, H., and Fritzsche, B. (2014). *Toward Translating Molecular Ear Development to Generate Hair Cells from Stem Cells, Adult Stem Cells*. New York, NY: Springer, 111–161.
- Zou, D., Silvius, D., Fritzsche, B., and Xu, P.-X. (2004). Eya1 and Six1 are essential for early steps of sensory neurogenesis in mammalian cranial placodes. *Development* 131, 5561–5572. doi: 10.1242/dev.01437

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 24 November 2014; accepted: 14 January 2015; published online: 05 February 2015.

Citation: Jahan I, Pan N and Fritzsche B (2015) Opportunities and limits of the one gene approach: the ability of Atoh1 to differentiate and maintain hair cells depends on the molecular context. *Front. Cell. Neurosci.* 9:26. doi: 10.3389/fncel.2015.00026

This article was submitted to the journal *Frontiers in Cellular Neuroscience*. Copyright © 2015 Jahan, Pan and Fritzsche. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.